



Review

Annual revision rates of partial versus total knee arthroplasty: A comparative meta-analysis



Harshvardhan Chawla^{a,b,*}, Jelle P. van der List^a, Alexander B. Christ^a, Maximiliano R. Sobrero^{a,b}, Hendrik A. Zuiderbaan^{a,c}, Andrew D. Pearle^a

^a Computer Assisted Surgery Center, Department of Orthopaedic Surgery, Hospital for Special Surgery, Weill Cornell Medical College, New York, NY, United States

^b Icahn School of Medicine at Mount Sinai, New York, NY, United States

^c Department of Orthopaedic Surgery, Spaarne Hospital, Hoofddorp, The Netherlands

ARTICLE INFO

Article history:

Received 17 July 2016

Received in revised form 29 October 2016

Accepted 7 November 2016

Keywords:

Annual revision rate

Total knee arthroplasty

Medial unicompartmental knee arthroplasty

Lateral unicompartmental knee arthroplasty

Patellofemoral arthroplasty

ABSTRACT

Background: Utilization of unicompartmental knee arthroplasty (UKA) and patellofemoral arthroplasty (PFA) as alternatives to total knee arthroplasty (TKA) for unicompartmental knee osteoarthritis (OA) has increased. However, no single resource consolidates survivorship data between TKA and partial resurfacing options for each variant of unicompartmental OA. This meta-analysis compared survivorship between TKA and medial UKA (MUKA), lateral UKA (LUKA) and PFA using annual revision rate as a standardized metric.

Methods: A systematic literature search was performed for studies quantifying TKA, MUKA, LUKA and/or PFA implant survivorship. Studies were classified by evidence level and assessed for bias using the MINORS and PEDro instruments. Annual revision rates were calculated for each arthroplasty procedure as percentages/observed component-year, based on a Poisson-normal model with random effects using the R-statistical software package.

Results: One hundred and twenty-four studies (113 cohort and 11 registry-based studies) met inclusion/exclusion criteria, providing data for 374,934 arthroplasties and 14,991 revisions. The overall evidence level was low, with 96.7% of studies classified as level III–IV. Annual revision rates were lowest for TKA (0.49%, CI 0.41 to 0.58), followed by MUKA (1.07%, CI 0.87 to 1.31), LUKA (1.13%, CI 0.69 to 1.83) and PFA (1.75%, CI 1.19 to 2.57). No difference was detected between revision rates for MUKA and LUKA ($p = 0.222$).

Conclusions: Revisions of MUKA, LUKA and PFA occur at an annual rate of 2.18, 2.31 and 3.57-fold that of TKA, respectively. These estimates may be used to inform clinical decision-making, guide patient expectations and evaluate the cost-effectiveness of total versus partial knee replacement in the setting of unicompartmental OA.

© 2016 Elsevier B.V. All rights reserved.

Contents

1.	Introduction	180
2.	Methods	181
2.1.	Search strategy	181
2.2.	Data extraction	181
2.3.	Statistical analysis	182

* Corresponding author at: Hospital for Special Surgery, 535 E. 70th Street, New York, NY 10021, United States.

E-mail addresses: chawlah@hss.edu (H. Chawla), vanderlistj@hss.edu (J.P. van der List), christa@hss.edu (A.B. Christ), maximiliano.sobrero@icahn.mssm.edu (M.R. Sobrero), hazuideraanmd@gmail.com (H.A. Zuiderbaan), pearlea@hss.edu (A.D. Pearle).

2.4. Heterogeneity and risk of bias assessment	182
3. Results	182
3.1. Search results	182
3.2. Annual revision rates	183
3.3. Heterogeneity and risk of bias	183
4. Discussion	183
5. Conclusions	186
Acknowledgments	186
References	186

1. Introduction

Partial knee arthroplasty (PKA) has undergone a renaissance over the past two decades for the treatment of unicompartmental knee osteoarthritis (OA) [1–3]. Based on the principle of selective resurfacing while sparing healthy bone and ligaments, PKA encompasses (1) medial unicompartmental knee arthroplasty (UKA), (2) lateral UKA and (3) patellofemoral arthroplasty (PFA) [2,4]. Short-term advantages of PKA include decreased blood loss, shorter operative times, lower complication rates and faster postoperative rehabilitation compared to total knee arthroplasty (TKA) [1,5–7]. In the long term, PKA provides greater range of motion (ROM) and higher rates of return to sports with preserved native kinematics and proprioception [2,5,6,8–13]. PKA has become increasingly popular as an alternative to TKA following improved survivorship reports (resulting from updated implant designs, surgical techniques and patient selection criteria) [2,4,14,15].

This represents a shift in the management philosophy of knee OA from a ‘one size fits all’ solution to a tricompartamental ‘hub and spoke’ model, in which PKA presents a joint-preserving option where indicated (Figure 1) [1–3,7]. To guide this clinical decision, a need exists for comparative survivorship data between TKA and corresponding partial resurfacing options for all variants




Figure 1. Compartmental approach to knee arthroplasty. Past versus present management approaches for isolated unicompartmental knee OA. Historically, the entire joint was replaced regardless of degenerative pattern (A). Under the modern tricompartamental model, only the diseased compartment is resurfaced for patients who qualify (B). Abbreviations: TKA, total knee arthroplasty; UKA, unicompartmental (tibiofemoral) arthroplasty; PFA, patellofemoral arthroplasty; OA, osteoarthritis.

of unicompartmental knee OA. To date, no single resource provides this information. Meta-analyses comparing UKA to TKA do not distinguish between medial and lateral subtypes, despite evidence that these comprise two very different procedures [16–21]. No meta-analysis or randomized clinical trials (RCT) comparing PFA and TKA survivorship exist [22]. Registries contain data for all such procedures but reflect the implants and techniques of a single nation, limiting generalizability. Combining findings from different sources is not possible, as survivorship is expressed non-uniformly with metrics ranging from cumulative failure to hazard ratios.

We performed a meta-analysis to create a comprehensive resource for survivorship data between TKA and each PKA alternative under the tricompartmental ‘hub and spoke’ model. Our objective was to permit direct comparison of annual revision rates between TKA and medial UKA (MUKA), lateral UKA (LUKA) or PFA for each variant of unicompartmental OA. This represents the first effort to consolidate survivorship data for all PKA subtypes against the existing standard of treatment.

2. Methods

2.1. Search strategy

A systematic literature search was performed in the PubMed, EMBASE and Cochrane Library databases on October 12, 2015 to identify studies reporting survivorship data for TKA, MUKA, LUKA and PFA. The search terms consisted of ‘(total OR TKA OR TKR),’ ‘(unicompartmental OR unicondylar OR UKA OR UKR),’ and ‘(patellofemoral OR PFA OR PFR)’ combined with ‘(knee arthroplasty OR knee replacement) AND (prosthesis OR implant) AND (failure OR revision OR reoperation OR survival OR survivorship)’ across three parallel searches (Figure 2). Results were filtered to retrieve only English-language studies published in 2005 or later. Two authors (HC and JPL) independently screened all entries by title and abstract against the inclusion/exclusion criteria (Table 1). Studies selected then underwent full-text review (HC and JPL) against these criteria. A third author (HAZ) was consulted in the event of any disagreement. Consensus was ultimately achieved regarding the inclusion/exclusion of all studies reviewed. The search and data analysis was performed in accordance with PRISMA guidelines [23].

2.2. Data extraction

Data describing (1) patient source (cohort or registry-based), (2) mean cohort age, (3) initial cohort size, (4) number of deaths with implant status unknown and/or losses to follow-up, (5) minimum follow-up duration, (6) mean follow-up duration, and (7) number of revisions was recorded in a spreadsheet (Excel 2008, Microsoft Inc., Redmond, WA) for all studies. Revision was defined as ‘a new operation in a previously resurfaced knee during which one or more of the components are exchanged, removed or added,’ per the Swedish Knee Registry [24]. Open reduction and internal fixation (ORIF) for periprosthetic fractures was counted towards this definition, as was subsequent patellar resurfacing.

Implant survivorship was reported as annual revision rate (expressed as percentage per observed component-year). This metric corrects for differential follow-up periods between populations, permitting direct comparison with a high degree of statistical accuracy [16,25]. All deaths with implant status unknown and losses to follow-up were subtracted from the initial population size,




Figure 2. Systematic search algorithm. Systematic search algorithm performed per PRISMA guidelines.

Table 1

Inclusion and exclusion criteria.

Inclusion	
1	Cohort or registry-based studies with TKA, UKA or PFA as index procedure
2	Primary osteoarthritis as the main indication for surgery
3	Reported mean postoperative follow-up time or total observed component-years
4	Reported total number and reasons for revision/reoperation
5	Minimum postoperative follow-up of two years
6	Minimum final population of 300 (TKA), 100 (medial UKA), 20 (lateral UKA) or 20 (PFA)
Exclusion	
1	Only included patients with specific pathology (i.e. hemophilia or morbid obesity) or demographic cutoff (i.e. only aged >65 or only female) not representative of the general population ^a
2	Only reported specific failure modes (e.g. revisions not “all-cause”)
3	Only reported revisions/reoperations for a single component
4	Previous bony surgeries (e.g. high tibial osteotomy, tibial tubercle transfer, patellectomy)
5	Revision or complex primary procedures (e.g. involving augmentation or bony defects)
6	Duplicate or overlapping patient populations
7	Did not report cohort size, loss to follow-up and/or revisions/reoperations separately for medial and lateral UKA if combined patient population

TKA, total knee arthroplasty; UKA, unicompartmental knee arthroplasty; PFA, patellofemoral arthroplasty. Where replicate patient data was detected (overlapping author/institution/registry, implant model and date range), the study with the greatest number of observed component-years was retained and redundant studies were excluded.

^a Exclusion criteria 1 was applied to papers that included only subjects with a specific pathology or demographic. This was done to ensure homogeneity between the cohorts being compared. Subjects with these features were still present throughout the studies selected for inclusion, but only if part of a representative population as opposed to an isolated special population.

which was based on the number of eligible subjects minus any exclusions, deaths, or losses prior to minimum follow-up. The resulting final cohort size was multiplied by mean follow-up to estimate the total observed component-years [26]. Annual revision rates were generated for each study by dividing the total number of revisions by total observed component-years. Three studies did not report mean follow-up but provided at least two of the following: (1) total revisions, (2) total observed component-years or (3) net annual revision rate [25,27,28]. These values were manually input and used to calculate the missing values.

2.3. Statistical analysis

The total number of knee arthroplasties, revisions and observed component-years were extracted for each procedure category, stratified by cohort only or combined cohort and registry-based studies. Pooled annual revision rates were calculated for (A) all studies and (B) cohort studies only using a Poisson-normal model with random effects. Log incidence rates were back-transformed using exponentiation to obtain annual revision rates for TKA, MUKA, LUKA and PFA. These were reported as mean values with 95% confidence intervals (CI) [29]. Significance testing was performed via meta-regression with a normal-normal mixed effects model using TKA as the reference value at a threshold of $p < 0.05$. Additional significance testing was performed between MUKA and LUKA annual revision rates. All statistical analysis was performed using R version 3.2.3 (R Foundation for Statistical Computing, Vienna, Austria) with Metafor version 1.9-8 (Maastricht University, Maastricht, Netherlands).

2.4. Heterogeneity and risk of bias assessment

All studies included were assessed for methodological, clinical and statistical heterogeneity. Methodological heterogeneity was evaluated concurrently with risk of bias by rating the evidence level and methodological quality of each publication, as this accounts for specific aspects of study design. This was done independently by two authors (HC and MRS), with any disagreements referred to a third author (JPL) and discussed until consensus was reached. Levels of evidence were based on the definitions published by the Center for Evidence-Based Medicine [30]. Quality assessment was performed with the PEDro (Physiotherapy Evidence Database) or MINORS (Methodological Index for Non-Randomized Studies) instruments. The former contains 11 items of which 10 are counted, and was applied to RCT [31]. The MINORS scale consists of eight items (with four supplementary items for comparative studies), and was used to assess non-randomized studies [32]. Clinical heterogeneity was evaluated by comparing the mean patient age between each arthroplasty cohort. This was reported as the mean and standard deviation weighted by final cohort size. Other patient variables were not compared due to inconsistency in how these were reported between studies. Statistical heterogeneity was assessed using the I^2 statistic with thresholds of 25% (low heterogeneity), 50% (moderate heterogeneity) and 75% (high heterogeneity).

3. Results

3.1. Search results

A total of 124 publications were selected for inclusion after full-text review (Figure 2). Three sources provided data for multiple procedures (two MUKA and LUKA, one TKA and PFA) [25,27,33], yielding 127 unique data entries in total. Fifty-two entries

Table 2
Study and population statistics.

	Mean age (Y)	Knees	Revisions	Component-years
<i>Cohort and registry-based</i>				
TKA	69.4 ± 1.6	333,727	12,907	2,058,368.7
Medial UKA	65.8 ± 2.7	38,742	1851	154,238.0
Lateral UKA	65.0 ± 5.0	968	73	5135.9
PFA	59.7 ± 5.7	1497	160	8412.1
<i>Cohort only</i>				
TKA	68.6 ± 2.2	54,777	2145	435,178.3
Medial UKA	67.4 ± 4.0	14,647	800	79,738.1
Lateral UKA	65.0 ± 5.0	968	73	5135.9
PFA	59.4 ± 6.4	1205	140	7282.9

TKA, total knee arthroplasty; UKA, unicompartmental knee arthroplasty; PFA, patellofemoral arthroplasty; Y, years.

were recorded for TKA [25,34–84], 37 for MUKA [27,28,33,85–118], 13 for LUKA [27,33,119–129] and 25 for PFA [25,130–153]. A total of 113 cohort studies (115 entries) [27,33–39,41–50,52–58,60–72,76,78,79,81–99,101–153] and 11 registry-based studies (12 entries) [25,28,40,51,59,73–75,77,80,100] were included.

3.2. Annual revision rates

The combined studies provided data for a total of 374,934 knee arthroplasties and 14,991 revisions. Subject age, total knees, total revisions and total component-years by procedure and study type are provided in Table 2. Combined cohort and registry-based data showed that annual revision rates were lowest for TKA (0.49%, CI 0.41 to 0.58), followed in order by MUKA (1.07%, CI 0.87 to 1.31), LUKA (1.13%, CI 0.69 to 1.83) and PFA (1.75%, CI 1.19 to 2.57) (Table 3) (Figures 3–5). No difference was detected between annual revision rates of MUKA and LUKA ($p = 0.222$). Following stratification to include only cohort-based data, a slight decrease was noted in the annual revision rates of TKA, MUKA and PFA (Table 4). Similarly, no significant difference was observed between MUKA and LUKA annual revision rates ($p = 0.217$).

3.3. Heterogeneity and risk of bias

One level I RCT was included [64]. Two publications were level II prospective studies [58,103]. Thirty-three studies [27,28,40,46–48,51,53,56,59,61,65,71,72,74,77,78,81,84,86,97,98,103,104,106,109,118,122,124,125,138,140,143] were level III retrospective studies and 87 [33–39,41–45,49,50,52,54,55,57,60,63,66–70,73,75,76,79,80,82,83,85,87–96,99–102,105,107,108,110–117,119–121,123,126–137,139,141,142,144–153] were level IV case series, with one registry report ineligible for grading [25]. The RCT was awarded seven points on the PEDro scale, satisfying 70% of the criteria [64]. Thirty-five non-randomized comparative (levels II and III) studies scored a mean of 18.5 Standard deviation (SD 2.4) on the MINORS instrument, corresponding to 77.1% of the maximum possible. Eighty-seven non-randomized non-comparative (level IV) studies scored a mean of 10.8 (SD 1.2) on the MINORS scale, representing 67.4% of the maximum points possible. Statistical heterogeneity was high among the TKA, MUKA and PFA studies and moderate to high for the LUKA studies in both the combined and cohort-only analyses (Tables 3–4).

4. Discussion

Our study synthesized clinical data from the past decade to compare implant survivorship for TKA versus PKA in all scenarios. Revision of MUKA (1.07%), LUKA (1.13%) and PFA (1.75%) was 2.18, 2.31 and 3.57-fold more frequent than revision of TKA (0.49%), respectively (Figures 3–5). This represents the first population-level comparison of such revision rates under the tricompartmental model, using a time-independent metric.

Several explanations exist for the lower survivorship of UKA compared to TKA. First, UKA is highly sensitive to technical parameters such as mechanical alignment [154,155]. Overcorrection induces degeneration of the contralateral compartment, leading to revision [156]. Second, definitions of aseptic loosening (which constitutes 25% of early failures) are inconsistent and user-dependent

Table 3
Comparative annual revision rates of cohort and registry-based studies.

	TKA	Medial UKA	Lateral UKA	PFA
Studies included	52	37	13	25
Annual revision rate	0.49	1.07	1.13	1.75
Confidence interval (95%)	0.41–0.58	0.87–1.31	0.69–1.83	1.19–2.57
<i>p</i> -Value	Reference	<0.001	<0.001	<0.001
I^2 (%)	98.67	92.50	65.97	79.89

TKA, total knee arthroplasty; UKA, unicompartmental knee arthroplasty; PFA, patellofemoral arthroplasty. Annual revision rate is expressed as the percentage of revisions per observed-component year. Significance testing was performed with meta-regression using TKA as the reference.




Figure 3. Predicted implant survivorship of TKA versus medial UKA. Projected cumulative implant survivorship of total knee arthroplasty (TKA) compared to medial unicompartmental knee arthroplasty (UKA) over an extrapolated period of 30 years. Transparent bands represent 95% confidence intervals.

[157]. Aseptic loosening is diagnosed by the presence of radiographic radiolucent lines (RLL) in the presence of postoperative pain [158]. Low-volume surgeons, however, may lack the experience to diagnose this phenomenon with specificity [158,159]. RLL are often physiologic, while residual pain can last for up to two years as tibial stresses redistribute [101]. Third, surgeons have a lower threshold to revise UKA due to the relative ease (versus primary TKA to revision TKA) and preserved bone stock [160]. Finally, OA progression comprises 40% of late UKA failures [157]. This reflects the natural history of a pathologic process rather than a mechanical shortcoming of the implant, affecting survivorship rates of UKA but not TKA.

Technological advances and policy initiatives may help UKA achieve survivorship parity with TKA. Robotic and navigated systems have shown promise in achieving narrow tolerances for key technical parameters [161,162]. However, long-term follow-up data is required. Preliminary outcomes of cementless UKA implants have also been encouraging, as these decrease the incidence of RLL that may prompt unnecessary revision in the presence of temporary pain [25,101]. At the policy level, several authors have endorsed the creation of regional UKA 'hubs' to decrease revision associated with low-volume facilities [163,164]. As implant durability continues to improve, UKA is likely to be increasingly regarded as a permanent rather than a temporizing alternative to TKA.

Notably, the data did not reveal significantly different annual revision rates between MUKA and LUKA. In recent years, research underscoring the kinematic and anatomic differences between the medial and lateral tibiofemoral compartments has led to these being regarded as two very distinct procedures [18–21]. Several authors have proposed that the lower volume and perception of LUKA as a more technically demanding procedure have contributed to its underutilization in favor of TKA [3,33,165]. A recent meta-analysis found no significant survivorship difference between MUKA and LUKA, with two cohort studies reaching similar conclusions [27,165,166]. Our study not only corroborates these findings with increased statistical power, but also provides a control baseline in the form of TKA revision rates obtained with identical methodology. This is particularly salient given the lack of any RCT comparing LUKA to TKA.

Higher revision rates for PFA may be explained by (1) non-uniform selection criteria, (2) implant design issues and (3) non-standardized surgical technique [7,15,167,168]. PFA lacks an equivalent to the 1989 Kozinn and Scott criteria, which decreased UKA revision rates through strict selection guidelines [2,169]. Tibiofemoral degeneration and/or malalignment are absolute contraindications for PFA given the risk of OA progression, with some authors suggesting that secondary OA (i.e. secondary to trochlear




Figure 4. Predicted implant survivorship of TKA versus lateral UKA. Projected cumulative implant survivorship of total knee arthroplasty (TKA) compared to lateral unicompartmental knee arthroplasty (UKA) over an extrapolated period of 30 years. Transparent bands represent 95% confidence intervals.




Figure 5. Predicted implant survivorship of TKA versus patellofemoral arthroplasty. Projected cumulative implant survivorship of total knee arthroplasty (TKA) compared to patellofemoral arthroplasty (PFA) over an extrapolated period of 30 years. Transparent bands represent 95% confidence intervals.

dysplasia, maltracking or trauma) represents a more suitable indication [135,147,167,170,171]. Gender-specific and customized implants are of particular interest in the development of third-generation PFA, with respect to minimizing maltracking and pain from soft-tissue impingement [150,172]. Intramedullary instrumentation (versus earlier freehand techniques) and robot-assisted techniques have also been shown to aid the precision and reproducibility of implant placement [140,150,172,173]. Second-generation PFA has already demonstrated a three to four-fold reduction in revision rates from earlier implants, rekindling interest in PFA as a means to delay TKA for young active patients [15,174].

The above findings have important implications for the surgical treatment of unicompartmental OA. Despite its high durability, TKA imposes significant functional and lifestyle restrictions secondary to the loss of ligaments and healthy bone stock [175,176]. Postoperative ROM is markedly better following PKA, particularly in flexion-dependent activities [5,9,10]. Subjects undergoing PKA report higher activity levels and return to sports versus those who receive TKA [6,11–13]. At the individual level, this is of particular importance to younger patients who may prefer PKA for its functional advantages despite the increased failure risk [2,174]. Annual revision rates permit direct comparison of this lifetime risk based on the age of the individual patient. At the policy level, the tradeoff between the clinical benefits of PKA against the higher risk of implant failure (with the associated costs and morbidity of revision TKA) determines the relative cost-effectiveness of such procedures [177,178]. As the US healthcare industry faces continuing pressure to demonstrate cost-efficiency, these time-independent failure rates may be applied to decision-analytic models quantifying the economics of selecting PKA over TKA [179].

This study has several limitations. First, no registry-based studies conforming to the inclusion/exclusion criteria provided survivorship data for LUKA [28]. As registry-based failure rates are typically higher than those of cohort studies, this may have the relative annual revision rates of LUKA [16]. However, subgroup analysis of cohort studies showed that this distinction had a negligible effect on the estimated rates for TKA, MUKA and PFA. Second, the overall quality of the studies selected was low with the majority of studies reporting level IV evidence. However, as no level I or II studies concerning LUKA or PFA have been published to date, the present study represents the best available effort to consolidate survivorship data across all four procedures [15,165,180]. Third, TKA studies were not matched to comparative PKA studies by indication (e.g. unicompartmental OA). This is unlikely to influence the findings or their clinical implications, however. PKA and TKA are generally subject to the same modes of failure, with the exception of multicompartmental OA progression [157,181]. The use of TKA obviates OA progression as an indication for revision, eliminating the ability to statistically differentiate between single or multicompartmental OA as the initial surgical indication. Finally, annual revision rates are known to inversely correlate with patient age at arthroplasty [24,25]. This relationship was observed in the present study, where patients undergoing PKA were generally younger. While younger age may predispose such patients to activity-dependent failure modes such as aseptic loosening, the most common mode of failure for all PKA variants remains OA progression [157,181]. This represents a primary etiology that manifests independent of patient age, partially mitigating this consideration. While age stratified survivorship statistics would have been informative, the scarcity of registry data and lack of age grouping in cohort

Table 4
Comparative annual revision rates of cohort studies.

	TKA	Medial UKA	Lateral UKA	PFA
Studies included	43	35	13	24
Annual revision rate	0.46	1.04	1.13	1.74
Confidence interval (95%)	0.37–0.57	0.84–1.30	0.69–1.83	1.16–2.61
p-Value	Reference	<0.001	<0.001	<0.001
I ² (%)	95.20	89.07	65.97	79.52

TKA, total knee arthroplasty; UKA, unicompartmental knee arthroplasty; PFA, patellofemoral arthroplasty. Annual revision rate is expressed as the percentage of revisions per observed-component year. Significance testing was performed with meta-regression using TKA as the reference.

studies did not permit such analysis. Future meta-analyses incorporating age in the search strategy will be required to further refine this survivorship data.

5. Conclusions

Annual revision rates of MUKA (1.07%), LUKA (1.13%) and PFA (2.75%) occurred at a ratio of 2.18, 2.31 and 3.57, respectively, versus that of TKA (0.49%). Furthermore, no significant differences were detected between the annual revision rates of MUKA and LUKA. These estimates may be used to inform clinical decision-making, guide patient expectations and/or evaluate the cost-effectiveness of total versus partial knee replacement in the setting of unicompartmental knee OA.

Acknowledgments

The authors wish to thank Lily Lee, MS and Kara Fields, MS from the Healthcare Research Institute at the Hospital for Special Surgery for their assistance with statistical analysis.

References

- [1] Bolognesi MP, Greiner MA, Attarian DE, Watters TS, Wellman SS, Curtis LH, et al. Unicompartmental knee arthroplasty and total knee arthroplasty among Medicare beneficiaries, 2000 to 2009. *J Bone Joint Surg Am* 2013;95, e174. <http://dx.doi.org/10.2106/JBJS.L.00652>.
- [2] Geller JA, Yoon RS, Macaulay W. Unicompartmental knee arthroplasty: a controversial history and a rationale for contemporary resurgence. *J Knee Surg* 2008; 21:7–14.
- [3] Parratte S, Ollivier M, Lunebourg A, Abdel MP, Argenson JN. Long-term results of compartmental arthroplasties of the knee: long term results of partial knee arthroplasty. *Bone Joint J* 2015;97B:9–15. <http://dx.doi.org/10.1302/0301-620X.97B10.36426>.
- [4] Blazina ME, Fox JM, Del Pizzo W, Broukhim B, Ivey FM. Patellofemoral replacement. *Clin Orthop Relat Res* 1979;98–102. <http://dx.doi.org/10.1097/01.blo.0000172301.66790.62>.
- [5] Yang KY, Wang MC, Yeo SJ, Lo NN. Minimally invasive unicondylar versus total condylar knee arthroplasty – early results of a matched-pair comparison. *Singapore Med J* 2003;44:559–62.
- [6] Dahm DL, Al-Rayashi W, Dajani K, Shah JP, Levy BA, Stuart MJ. Patellofemoral arthroplasty versus total knee arthroplasty in patients with isolated patellofemoral osteoarthritis. *Am J Orthop (Belle Mead NJ)* 2010;39:487–91.
- [7] Ackroyd CE, Chir B. Development and early results of a new patellofemoral arthroplasty. *Clin Orthop Relat Res* 2005;7–13. <http://dx.doi.org/10.1097/01.blo.0000171914.94503.d1>.
- [8] van Jonbergen HPW, Werkman DM, Barnaart LF, van Kampen A. Long-term outcomes of patellofemoral arthroplasty. *J Arthroplasty* 2010;25:1066–71. <http://dx.doi.org/10.1016/j.arth.2009.08.023>.
- [9] Wiik AV, Aqil A, Tankard S, Amis AA, Cobb JP. Downhill walking gait pattern discriminates between types of knee arthroplasty: improved physiological knee functionality in UKA versus TKA. *Knee Surg Sports Traumatol Arthrosc* 2015;23:1748–55. <http://dx.doi.org/10.1007/s00167-014-3240-x>.
- [10] Horikawa A, Miyakoshi N, Shimada Y, Kodama H. Comparison of clinical outcomes between total knee arthroplasty and unicompartmental knee arthroplasty for osteoarthritis of the knee: a retrospective analysis of preoperative and postoperative results. *J Orthop Surg Res* 2015;10:168. <http://dx.doi.org/10.1186/s13018-015-0309-2>.
- [11] Walton NP, Jahromi I, Lewis PL, Dobson PJ, Angel KR, Campbell DG. Patient-perceived outcomes and return to sport and work: TKA versus mini-incision unicompartmental knee arthroplasty. *J Knee Surg* 2006;19:112–6.
- [12] Hopper GP, Leach WJ. Participation in sporting activities following knee replacement: total versus unicompartmental. *Knee Surg Sports Traumatol Arthrosc* 2008;16:973–9. <http://dx.doi.org/10.1007/s00167-008-0596-9>.
- [13] Ho JC, Stitzlein RN, Green CJ, Stoner T, Froimson MI. Return to sports activity following UKA and TKA. *J Knee Surg* 2015. <http://dx.doi.org/10.1055/s-0035-1551835>.
- [14] Insall J, Aglietti P. A five to seven-year follow-up of unicondylar arthroplasty. *J Bone Joint Surg Am* 1980;62:1329–37.
- [15] Dy CJ, Franco N, Ma Y, Mazumdar M, McCarthy MM, Gonzalez Della Valle A. Complications after patello-femoral versus total knee replacement in the treatment of isolated patello-femoral osteoarthritis. A meta-analysis. *Knee Surg Sports Traumatol Arthrosc* 2012;20:2174–90. <http://dx.doi.org/10.1007/s00167-011-1677-8>.
- [16] Pabinger C, Berghold A, Boehler N, Labek G. Revision rates after knee replacement: cumulative results from worldwide clinical studies versus joint registers. *Osteoarthritis Cartilage* 2013;21:263–8. <http://dx.doi.org/10.1016/j.joca.2012.11.014>.
- [17] Arirachakaran A, Choovit P, Putananon C, Muangsiin S, Kongtharvonskul J. Is unicompartmental knee arthroplasty (UKA) superior to total knee arthroplasty (TKA)? A systematic review and meta-analysis of randomized controlled trial. *Eur J Orthop Surg Traumatol Orthop Traumatol* 2015;25:799–806. <http://dx.doi.org/10.1007/s00590-015-1610-9>.
- [18] Weidow J, Mars I, Kärrholm J. Medial and lateral osteoarthritis of the knee is related to variations of hip and pelvic anatomy. *Osteoarthritis Cartilage* 2005;13: 471–7. <http://dx.doi.org/10.1016/j.joca.2005.01.009>.
- [19] Weidow J, Pak J, Kärrholm J, Weidow J, Pak J, Kärrholm J. Different patterns of cartilage wear in medial and lateral gonarthrosis. 6470; 2016. <http://dx.doi.org/10.1080/000164702320155347>.
- [20] Servien E, Saffarini M, Lustig S, Chomel S, Neyret P. Lateral versus medial tibial plateau: morphometric analysis and adaptability with current tibial component design. *Knee Surg Sports Traumatol Arthrosc* 2008;16:1141–5. <http://dx.doi.org/10.1007/s00167-008-0620-0>.
- [21] Kaya Bicer E, Servien E, Lustig S, Demey G, Ait Si Selmi T, Neyret P. Sagittal flexion angle of the femoral component in unicompartmental knee arthroplasty: is it same for both medial and lateral UKAs? *Knee Surg Sports Traumatol Arthrosc* 2010;18:928–33. <http://dx.doi.org/10.1007/s00167-010-1063-y>.
- [22] Odumenya M, McGuinness K, Achten J, Parsons N, Spalding T, Costa M. The Warwick patellofemoral arthroplasty trial: a randomised clinical trial of total knee arthroplasty versus patellofemoral arthroplasty in patients with severe arthritis of the patellofemoral joint. *BMC Musculoskelet Disord* 2011;12:265. <http://dx.doi.org/10.1186/1471-2474-12-265>.
- [23] Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol* 2009;62:e1–34. <http://dx.doi.org/10.1016/j.jclinepi.2009.06.006>.
- [24] Robertsson O, W-Dahl A. Swedish knee arthroplasty register: annual report. Lund; 2015.
- [25] Rothwell A, Hobbs T, Frampton C, Armour P, Muir D, Mohammed K. The New Zealand Joint Registry: fifteen year report, January 1999 to December 2013; 2014.
- [26] van der List JP, Chawla H, Zuiderbaan HA, Pearle AD. The role of preoperative patient characteristics on outcomes of unicompartmental knee arthritis: a meta-analysis critique. *J Arthroplasty* 2016;1–23. <http://dx.doi.org/10.1016/j.arth.2016.04.001>.
- [27] Liebs TR, Herzberg W. Better quality of life after medial versus lateral unicondylar knee arthroplasty knee. *Clin Orthop Relat Res* 2013;471:2629–40. <http://dx.doi.org/10.1007/s11999-013-2966-y>.
- [28] Baker P, Jameson S, Critchley R, Reed M, Gregg P, Deehan D. Center and surgeon volume influence the revision rate following unicondylar knee replacement: an analysis of 23,400 medial cemented unicondylar knee replacements. *J Bone Joint Surg Am* 2013;95:702–9. <http://dx.doi.org/10.2106/JBJS.L.00520>.
- [29] Niël-Weise BS, Stijnen T, van den Broek PJ. Anti-infective-treated central venous catheters for total parenteral nutrition or chemotherapy: a systematic review. *J Hosp Infect* 2008;69:114–23. <http://dx.doi.org/10.1016/j.jhin.2008.02.020>.

- [30] Levels of evidence. Oxford Cent Evidence-Based Med; 2009.
- [31] Morton NAD. The PEDro scale is a valid measure of the methodological quality of clinical trials: a demographic study. *Aust J Physiother* 2009;55:129–33. [http://dx.doi.org/10.1016/S0004-9514\(09\)70043-1](http://dx.doi.org/10.1016/S0004-9514(09)70043-1).
- [32] Slim K, Nini E, Forestier D, Kwiatkowski F, Panis Y, Chipponi J. Methodological index for non-randomized studies (minors): development and validation of a new instrument. *ANZ J Surg* 2003;73:712–6.
- [33] Xing Z, Ph D, Katz J, Jiranek W. Unicompartmental knee arthroplasty: factors in fluencing the outcome, 1; 2012.
- [34] Argenson JN, Boisgard S, Parratte S, Descamps S, Bercovy M, Bonnevalle P, et al. Survival analysis of total knee arthroplasty at a minimum 10 years' follow-up: a multicenter French nationwide study including 846 cases. *Orthop Traumatol Surg Res* 2013;99:385–90. <http://dx.doi.org/10.1016/j.otsr.2013.03.014>.
- [35] Attar FG, Khaw FM, Kirk LMG, Gregg PJ. Survivorship analysis at 15 years of cemented press-fit condylar total knee arthroplasty. *J Arthroplasty* 2008;23:344–9. <http://dx.doi.org/10.1016/j.arth.2007.02.012>.
- [36] Bae DK, Song SJ, Park MJ, Eoh JH, Song JH, Park CH. Twenty-year survival analysis in total knee arthroplasty by a single surgeon. *J Arthroplast* 2012;27:1297–304.e1. <http://dx.doi.org/10.1016/j.arth.2011.10.027>.
- [37] Bauze AJ, Falworth MS, Oakeshott RD. Early results of total knee arthroplasty with a low contact stress anteroposterior glide. *J Orthop Surg (Hong Kong)* 2009;17:174–8.
- [38] Bercovy M, Langlois J, Beldame J, Lefebvre B. Functional results of the ROCC mobile bearing knee. 602 cases at midterm follow-up (5 to 14 years). *J Arthroplasty* 2014;30:973–9. <http://dx.doi.org/10.1016/j.arth.2015.01.003>.
- [39] Bistolfi A, Lee GC, Deledda D, Rosso F, Berchiolla P, Crova M, et al. NexGen LPS mobile bearing total knee arthroplasty: 10-year results. *Knee Surg Sports Traumatol Arthrosc* 2014;22:1786–92. <http://dx.doi.org/10.1007/s00167-014-3019-0>.
- [40] Bordini B, Stea S, Cremonini S, Viceconti M, De Palma R, Toni A. Relationship between obesity and early failure of total knee prostheses. *BMC Musculoskelet Disord* 2009;10:29. <http://dx.doi.org/10.1186/1471-2474-10-29>.
- [41] Borriero F, Bonnevalle P, Mabit C, Guingand O, Bertin D, Bonnomet F, et al. Scorpio single radius total knee arthroplasty. A minimal five-year follow-up multicentric study. *Int Orthop* 2011;35:1777–82. <http://dx.doi.org/10.1007/s00264-011-1226-6>.
- [42] Chalidis BE, Sachinis NP, Papadopoulos P, Petsatodis E, Christodoulou AG, Petsatodis G. Long-term results of posterior-cruciate-retaining Genesis i total knee arthroplasty. *J Orthop Sci* 2011;16:726–31. <http://dx.doi.org/10.1007/s00776-011-0152-1>.
- [43] Chatain F, Gaillard TH, Denjean S, Tayot O. Outcomes of 447SCORE highly congruent mobile-bearing total knee arthroplasties after 5–10 years follow-up. *Orthop Traumatol Surg Res* 2013;99:681–6. <http://dx.doi.org/10.1016/j.otsr.2013.05.003>.
- [44] Chiavetta J, Fehring TK, Odum S, Griffin W, Mason JB. Importance of a balanced-gap technique in rotating-platform knees. *Orthopedics* 2006;29:S45–8.
- [45] Cho WS, Youm YS, Ahn SC, Sohn DW. What have we learned from LCS mobile-bearing knee system? *Knee Surg Sports Traumatol Arthrosc* 2010;18:1345–51. <http://dx.doi.org/10.1007/s00167-010-1166-5>.
- [46] Dalury DF, Mason JB, Murphy JA, Adams MJ. Analysis of the outcome in male and female patients using a unisex total knee replacement system. *J Bone Joint Surg Br* 2009;91:357–60. <http://dx.doi.org/10.1302/0301-620X.91B3.21771>.
- [47] Demey G, Hobbs H, Lustig S, Servien E, Trouillet F, Magnusson RA, et al. Influence of gender on the outcome of total knee arthroplasty. *Eur Orthop Traumatol* 2012;3:11–6. <http://dx.doi.org/10.1007/s12570-012-0094-x>.
- [48] Deshmukh AJ, Rathod PA, Moses MJ, Snir N, Marwin SE, Dayan AJ. Does a non-stemmed constrained condylar prosthesis predispose to early failure of primary total knee arthroplasty? *Knee Surg Sports Traumatol Arthrosc* 2015;11:3. <http://dx.doi.org/10.1007/s00167-014-3494-3>.
- [49] Efstathopoulos N, Mavrogenis AF, Lallou S, Nikolaou V, Papagelopoulos PJ, Savvidou OD, et al. 10-year evaluation of the cementless low-contact-stress rotating-platform total knee arthroplasty. *J Long Term Eff Med Implants* 2009;19:255–63.
- [50] García-Crespo R, Marco F, Moro LE, Ariza G, Luque R, López-Durán L. Midterm results of Optetrak posterior-stabilized total knee system after 7 to 12 years in a university hospital. *J Arthroplasty* 2011;26:1326–31. <http://dx.doi.org/10.1016/j.arth.2010.11.020>.
- [51] Goe TJ, Sharma A, Tatman P, Mehle S. Do "premium" joint implants add value?: analysis of high cost joint implants in a community registry. *Clin Orthop Relat Res* 2011;469:48–54. <http://dx.doi.org/10.1007/s11999-010-1436-z>.
- [52] Harwin SF, Elmallah RK, Jauregui JJ, Cherian JJ, Mont MA. Outcomes of a newer-generation cementless total knee arthroplasty design. *Orthopedics* 2015;38:620–4. <http://dx.doi.org/10.3928/01477447-20151002-04>.
- [53] Hickey BA, Kempshall PJ, Metcalfe AJ, Forster MC. Review of Kinemax total knee replacements performed at the NHS treatment centre, Weston-Super-Mare: five-year results. *Bone Joint J* 2012;94-B:510–6. <http://dx.doi.org/10.1302/0301-620X.94B4.27978>.
- [54] Hosaka K, Saito S, Ishii T, Mori S, Sumino T, Tokuhashi Y. Asian-Specific total knee system: 5–14 year follow-up study. *BMC Musculoskelet Disord* 2011;12:251. <http://dx.doi.org/10.1186/1471-2474-12-251>.
- [55] Hunter N, Clayton RAE, Brenkel JJ. Press fit condylar Sigma total knee arthroplasty: 7–9 years results. *Eur J Orthop Surg Traumatol* 2009;19:409–13. <http://dx.doi.org/10.1007/s00590-009-0446-6>.
- [56] Jackson MP, Sexton SA, Yeung E, Walter WL, Walter WK, Zicat BA. The effect of obesity on the mid-term survival and clinical outcome of cementless total hip replacement. *J Bone Joint Surg Br* 2009;91:1296–300. <http://dx.doi.org/10.1302/0301-620X.91B10.22544>.
- [57] Kievit AJ, Schaafroth MU, Blankevoort L, Siersevelt IN, van Dijk CN, van Geenen RCI. Early experience with the vanguard complete total knee system: 2–7 years of follow-up and risk factors for revision. *J Arthroplasty* 2014;29:348–54. <http://dx.doi.org/10.1016/j.arth.2013.05.018>.
- [58] Kim Y-H, Kwon O-R, Kim J-S. Is one-stage bilateral sequential total hip replacement as safe as unilateral total hip replacement? *J Bone Joint Surg Br* 2009;91:316–20. <http://dx.doi.org/10.1302/0301-620X.91B11>.
- [59] Kremers HM, Sierra RJ, Schleck CD, Berry DJ, Cabanela ME, Hanssen AD, et al. Comparative survivorship of different tibial designs. *J Bone Joint Surg* 2014;121:1–7. <http://dx.doi.org/10.2106/JBJS.M.00820>.
- [60] Lizaur-Utrilla A, Gonzalez-Parreño S, Miralles-Muñoz FA, Lopez-Prats FA, Gil-Guillen V. Patient-related predictors of treatment failure after primary total knee arthroplasty for osteoarthritis. *J Arthroplasty* 2014;29:2095–9. <http://dx.doi.org/10.1016/j.arth.2014.07.011>.
- [61] Lombardi AV, Berend KR, Adams JB. Minimally invasive surgery is not a risk factor of early total knee arthroplasty failure. *Semin Arthroplasty* 2010;21:92–6. <http://dx.doi.org/10.1053/j.sart.2009.12.016>.
- [62] MacDonald SJ, Charron KD, Bourne RB, Naudie DD, McCalden RW, Rorabeck CH. The John Insall award: gender-specific total knee replacement: prospectively collected clinical outcomes. *Clin Orthop Relat Res* 2008;466:2612–6. <http://dx.doi.org/10.1007/s11999-008-0430-1>.
- [63] Mahoney OM, Kinsey TL. 5- to 9-year survivorship of single-radius, posterior-stabilized TKA. *Clin Orthop Relat Res* 2008;466:436–42. <http://dx.doi.org/10.1007/s11999-007-0070-x>.
- [64] Mahoney OM, Kinsey TL, D'Errico TJ, Shen J. The John Insall Award: no functional advantage of a mobile bearing posterior stabilized TKA. *Clin Orthop Relat Res* 2012;470:33–44. <http://dx.doi.org/10.1007/s11999-011-2114-5>.
- [65] Meding JB, Berend ME, Ritter MA, Galley MR, Malinzak RA. Intramedullary vs extramedullary femoral alignment guides. A 15-year follow-up of survivorship. *J Arthroplasty* 2011;26:591–5. <http://dx.doi.org/10.1016/j.arth.2010.05.008>.
- [66] Melton JTK, Mayahi R, Baxter SE, Facek M, Glezoc C. Long-term outcome in an uncemented, hydroxyapatite-coated total knee replacement: a 15- to 18-year survivorship analysis. *J Bone Joint Surg Br* 2012;94:1067–70. <http://dx.doi.org/10.1302/0301-620X.94B8.28350>.
- [67] Metsovitis SR. Rotaglide total knee arthroplasty: a long-term follow-up study. *J Bone Joint Surg* 2011;93:878. <http://dx.doi.org/10.2106/JBJS.I.01702>.
- [68] Mohammed A, Metcalfe A, Woodnutt D. Medium-term outcome of titanium nitride, mobile bearing total knee replacement. *Acta Orthop Belg* 2014;80:269–75.
- [69] Munzinger UK, Maffuletti NA, Guggi T, Bizzini M, Preiss S, Drobny T. Five-year results of the Innex total knee arthroplasty system. *Int Orthop* 2010;34:1159–65. <http://dx.doi.org/10.1007/s00264-009-0887-x>.
- [70] Nakamura S, Kobayashi M, Ito H, Nakamura K, Ueo T, Nakamura T. The bi-surface total knee arthroplasty: minimum 10-year follow-up study. *Knee* 2010;17:274–8. <http://dx.doi.org/10.1016/j.knee.2010.02.015>.
- [71] Nam D, Umunna BPN, Cross MB, Reinhardt KR, Duggal S, Cornell CN. Clinical results and failure mechanisms of a nonmodular constrained knee without stem extensions. *HSS J* 2012;8:96–102. <http://dx.doi.org/10.1007/s11420-012-9277-9>.

- [72] Nassif JM, Pietrzak WS. Clinical outcomes in men and women following total knee arthroplasty with a high-flex knee: no clinical effect of gender. *Sci World J* 2015;2015:1–6. <http://dx.doi.org/10.1155/2015/285919>.
- [73] Niemeläinen M, Skyttä ET, Remes V, Mäkelä K, Eskelinen A. Total knee arthroplasty with an uncemented trabecular metal tibial component: a registry-based analysis. *J Arthroplasty* 2014;29:57–60. <http://dx.doi.org/10.1016/j.arth.2013.04.014>.
- [74] Niinimäki T, Eskelinen A, Mäkelä K, Ohtonen P, Puhto AP, Remes V. Unicompartmental knee arthroplasty survivorship is lower than TKA survivorship: a 27-year Finnish registry study. *Clin Orthop Relat Res* 2014;472:1496–501. <http://dx.doi.org/10.1007/s11999-013-3347-2>.
- [75] Paxton EW, Inacio MCS, Khatod M, Yue E, Funahashi T, Barber T. Risk calculators predict failures of knee and hip arthroplasties: findings from a large health maintenance organization. *Clin Orthop Relat Res* 2015;473:3965–73. <http://dx.doi.org/10.1007/s11999-015-4506-4>.
- [76] Pelt CE, Erickson J, Christensen BA, Widmer B, Severson EP, Evans D, et al. The use of a modular titanium baseplate with a press-fit keel implanted with a surface cementing technique for primary total knee arthroplasty. *Biomed Res Int* 2014;2014. <http://dx.doi.org/10.1155/2014/972615>.
- [77] Petursson G, Fenstad AM, Havelin LI, Gøthesen Ø, Håkon S, Lygre L, et al. Better survival of hybrid total knee arthroplasty compared to cemented arthroplasty. *3674*; 2015. <http://dx.doi.org/10.3109/17453674.2015.1073539>.
- [78] Piedade SR, Pinaroli A, Servien E, Neyret P. Is previous knee arthroscopy related to worse results in primary total knee arthroplasty? *Knee Surg Sports Traumatol Arthrosc* 2009;17:328–33. <http://dx.doi.org/10.1007/s00167-008-0669-9>.
- [79] Roberts VI, Esler CNA, Harper WM. A 15-year follow-up study of 4606 primary total knee replacements. *J Bone Joint Surg Br* 2007;89:1452–6. <http://dx.doi.org/10.1302/0301-620X.89B11.19783>.
- [80] Robertsson O, Ranstam J, Lidgren L. Variation in outcome and ranking of hospitals: an analysis from the Swedish knee arthroplasty register. *Acta Orthop* 2006;77:487–93. <http://dx.doi.org/10.1080/17453670610046442>.
- [81] Schnurr C, Güdden I, Eysel P, König DP. Influence of computer navigation on TKA revision rates. *Int Orthop* 2012;36:2255–60. <http://dx.doi.org/10.1007/s00264-012-1606-6>.
- [82] Schroer WC, Stormont DM, Pietrzak WS. Seven-year survivorship and functional outcomes of the high-flexion vanguard complete knee system. *J Arthroplasty* 2014;29:61–5. <http://dx.doi.org/10.1016/j.arth.2013.04.018>.
- [83] Scott CEH, Clement ND, MacDonald DJ, Hamilton DF, Gaston P, Howie CR, et al. Five-year survivorship and patient-reported outcome of the Triathlon single-radius total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 2014;23:1676–83. <http://dx.doi.org/10.1007/s00167-014-2922-8>.
- [84] Stiehl JB, Hamelnyck KJ, Voorhorst PE. International multi-centre survivorship analysis of mobile bearing total knee arthroplasty. *Int Orthop* 2006;30:190–9. <http://dx.doi.org/10.1007/s00264-005-0053-z>.
- [85] Baur J, Zwicky L, Hirschmann MT, Ilchmann T, Clauss M. Metal backed fixed-bearing unicondylar knee arthroplasties using minimal invasive surgery: a promising outcome analysis of 132 cases. *BMC Musculoskelet Disord* 2015;16:177. <http://dx.doi.org/10.1186/s12891-015-0651-x>.
- [86] Bhattacharya R, Scott CEH, Morris HE, Wade F, Nutton RW. Survivorship and patient satisfaction of a fixed bearing unicompartmental knee arthroplasty incorporating an all-polyethylene tibial component. *Knee* 2012;19:348–51. <http://dx.doi.org/10.1016/j.knee.2011.04.009>.
- [87] Bruni D, Gagliardi M, Akkawi I, Raspugli GF, Bignozzi S, Marko T, et al. Good survivorship of all-polyethylene tibial component UKA at long-term follow-up. *Knee Surg Sports Traumatol Arthrosc* 2014. <http://dx.doi.org/10.1007/s00167-014-3361-2>.
- [88] Burnett RSJ, Nair R, Hall CA, Jacks DA, Pugh L, McAllister MM. Results of the Oxford Phase 3 mobile bearing medial unicompartmental knee arthroplasty from an independent center: 467 knees at a mean 6-year follow-up: analysis of predictors of failure. *J Arthroplasty* 2014;29:193–200. <http://dx.doi.org/10.1016/j.arth.2014.01.035>.
- [89] Campbell D, Schuster AJ, Pfluger D, Hoffmann F. Unicondylar knee replacement with a new tensioner device: clinical results of a multicentre study on 168 cases. *Arch Orthop Trauma Surg* 2010;130:727–32. <http://dx.doi.org/10.1007/s00402-010-1092-5>.
- [90] Chatellard R, Sauleau V, Colmar M, Robert H, Raynaud G, Brillhault J. Medial unicompartmental knee arthroplasty: does tibial component position influence clinical outcomes and arthroplasty survival? *Orthop Traumatol Surg Res* 2013;99:S219–25. <http://dx.doi.org/10.1016/j.otsr.2013.03.004>.
- [91] Choy WS, Kim KJ, Lee SK, Yang DS, Lee NK. Mid-term results of oxford medial unicompartmental knee arthroplasty. *Clin Orthop Surg* 2011;3:178–83. <http://dx.doi.org/10.4055/cios.2011.3.3.178>.
- [92] Clark M, Campbell DG, Kiss G, Dobson PJ, Lewis PL. Reintervention after mobile-bearing Oxford unicompartmental knee arthroplasty. *Clin Orthop Relat Res* 2010;468:576–80. <http://dx.doi.org/10.1007/s11999-009-1089-y>.
- [93] Dervin GF, Carruthers C, Feibel RJ, Giachino AA, Kim PR, Thurston PR. Initial experience with the Oxford unicompartmental knee arthroplasty. *J Arthroplasty* 2011;26:192–7. <http://dx.doi.org/10.1016/j.arth.2010.02.007>.
- [94] Edmondson M, Atrey A, East D, Ellens N, Miles K, Goddard R, et al. Survival analysis and functional outcome of the Oxford unicompartmental knee replacement up to 11 years follow up at a District General Hospital. *J Orthop* 2015;12:S105–10. <http://dx.doi.org/10.1016/j.jor.2013.12.007>.
- [95] Eickmann TH, Collier MB, Sukezaki F, McAuley JP, Engh GA. Survival of medial unicondylar arthroplasties placed by one surgeon 1984–1998. *Clin Orthop Relat Res* 2006;452:143–9. <http://dx.doi.org/10.1097/01.blo.0000238793.74843.dc>.
- [96] Hamilton WG, Ammeen DJ, Hopper RH. Mid-term survivorship of minimally invasive unicompartmental arthroplasty with a fixed-bearing implant: revision rate and mechanisms of failure. *J Arthroplasty* 2014;29:989–92. <http://dx.doi.org/10.1016/j.arth.2013.10.010>.
- [97] Heyse TJ, Efe T, Rumpf S, Schofer MD, Fuchs-Winkelmann S, Schmitt J, et al. Minimally invasive versus conventional unicompartmental knee arthroplasty. *Arch Orthop Trauma Surg* 2011;131:1287–90. <http://dx.doi.org/10.1007/s00402-011-1274-9>.
- [98] Hurst JM, Berend KR, Morris MJ, Lombardi AV. Abnormal preoperative MRI does not correlate with failure of UKA. *J Arthroplasty* 2013;28:184–6. <http://dx.doi.org/10.1016/j.arth.2013.05.011>.
- [99] Ji JH, Park SE, Song IS, Kang H, Ha JY, Jeong JJ. Complications of medial unicompartmental knee arthroplasty. *CIOS Clin Orthop Surg* 2014;6:365–72. <http://dx.doi.org/10.4055/cios.2014.6.4.365>.
- [100] Kristensen PW, Holm HA, Varnum C. Up to 10-year follow-up of the Oxford medial partial knee arthroplasty – 695 cases from a single institution. *J Arthroplasty* 2013;28:195–8. <http://dx.doi.org/10.1016/j.arth.2013.05.010>.
- [101] Liddle AD, Pit H, O'Brien S, Doran E, Penny ID, Hooper GJ, et al. Cementless fixation in Oxford unicompartmental knee replacement: a multicentre study of 1000 knees. *Bone Joint J* 2013;95B:181–7. <http://dx.doi.org/10.1302/0301-620X.95B2>.
- [102] Lim H-C, Bae J-H, Song S-H, Kim S-J. Oxford phase 3 unicompartmental knee replacement in Korean patients. *J Bone Joint Surg Br* 2012;94:1071–6. <http://dx.doi.org/10.1302/0301-620X.94B8.29372>.
- [103] Lyons MC, MacDonald SJ, Somerville LE, Naudie DD, McCalden RW. Unicompartmental versus total knee arthroplasty database analysis: is there a winner? *Clin Orthop Relat Res* 2012;470:84–90. <http://dx.doi.org/10.1007/s11999-011-2144-z>.
- [104] Maier MW, Kuhs F, Streit MR, Schuhmacher P, Walker T, Ewerbeck V, et al. Unicompartmental knee arthroplasty in patients with full versus partial thickness cartilage loss (PTCL): equal in clinical outcome but with higher reoperation rate for patients with PTCL. *Arch Orthop Trauma Surg* 2015;135:1169–75. <http://dx.doi.org/10.1007/s00402-015-2236-4>.
- [105] Matharu G, Robb C, Baloch K, Pynsent P. The Oxford medial unicompartmental knee replacement: survival and the affect of age and gender. *Knee* 2012;19:913–7. <http://dx.doi.org/10.1016/j.knee.2012.03.004>.
- [106] Murray DW, Pandit H, Weston-Simons JS, Jenkins C, Gill HS, Lombardi AV, et al. Does body mass index affect the outcome of unicompartmental knee replacement? *Knee* 2013;20:461–5. <http://dx.doi.org/10.1016/j.knee.2012.09.017>.
- [107] O'Donnell T, Neil MJ. The Repicci II unicondylar knee arthroplasty: 9-year survivorship and function. *Clin Orthop Relat Res* 2010;468:3094–102. <http://dx.doi.org/10.1007/s11999-010-1474-6>.
- [108] O'Rourke MR, Gardner JJ, Callaghan JJ, Liu SS, Goetz DD, Vittetoe DA, et al. The John Insall Award: unicompartmental knee replacement. *Clin Orthop Relat Res* 2005;440:27–37. <http://dx.doi.org/10.1097/01.blo.0000185451.96987.aa>.
- [109] Parratte S, Pauly V, Aubaniac JM, Argenson JNA. No long-term difference between fixed and mobile medial unicompartmental arthroplasty. *Clin Orthop Relat Res* 2012;470:61–8. <http://dx.doi.org/10.1007/s11999-011-1961-4>.
- [110] Price AJ, Orth F, Waite JC, Svard U. Long-term clinical results of the medial Oxford unicompartmental knee arthroplasty. *Clin Orthop Relat Res* 2005;171–80. <http://dx.doi.org/10.1097/01.blo.0000155347.89533.c5>.

- [111] Saenz CL, McGrath MS, Marker DR, Seyler TM, Mont MA, Bonutti PM. Early failure of a unicompartmental knee arthroplasty design with an all-polyethylene tibial component. *Knee* 2010;17:53–6. <http://dx.doi.org/10.1016/j.knee.2009.05.007>.
- [112] Schlueter-Brust K, Kugland K, Stein G, Henckel J, Christ H, Eysel P, et al. Ten year survivorship after cemented and uncemented medial Uniglide unicompartmental knee arthroplasties. *Knee* 2014;21:964–70. <http://dx.doi.org/10.1016/j.knee.2014.03.009>.
- [113] Smith TO, Clark A, Glasgow MMS, Donell ST. The mid-term clinical results of the phase 3 Oxford unicompartmental knee arthroplasty: a 6- to 8-year follow-up. *Eur J Orthop Surg Traumatol* 2012;22:307–14. <http://dx.doi.org/10.1007/s00590-011-0828-4>.
- [114] Steele RG, Hutabarat S, Evans RL, Ackroyd CE, Newman JH. Survivorship of the St George SLED medial unicompartmental knee replacement beyond ten years. *J Bone Joint Surg Br* 2006;88:1164–8. <http://dx.doi.org/10.1302/0301-620X.88B9.18044>.
- [115] Vasso M, Del Regno C, Perisano C, D'Amelio A, Corona K, Schiavone PA. Unicompartmental knee arthroplasty is effective: ten year results. *Int Orthop* 2015. <http://dx.doi.org/10.1007/s00264-015-2809-4>.
- [116] White SH, Roberts S, Kuiper JH. The cemented twin-peg Oxford partial knee replacement survivorship: a cohort study. *Knee* 2014;22:333–7. <http://dx.doi.org/10.1016/j.knee.2015.03.011>.
- [117] Yoshida K, Tada M, Yoshida H, Takei S, Fukuoka S, Nakamura H. Oxford phase 3 unicompartmental knee arthroplasty in Japan – clinical results in greater than one thousand cases over ten years. *J Arthroplasty* 2013;28:168–71. <http://dx.doi.org/10.1016/j.arth.2013.08.019>.
- [118] Zambianchi F, Digennaro V, Giorgini A, Grandi G, Fiacchi F, Mugnai R, et al. Surgeon's experience influences UKA survivorship: a comparative study between all-poly and metal back designs. *Knee Surg Sports Traumatol Arthrosc* 2014;23:2074–80. <http://dx.doi.org/10.1007/s00167-014-2958-9>.
- [119] Altuntas AO, Alsop H, Cobb JP. Early results of a domed tibia, mobile bearing lateral unicompartmental knee arthroplasty from an independent centre. *Knee* 2013;20:466–70. <http://dx.doi.org/10.1016/j.knee.2012.11.008>.
- [120] Argenson JNA, Parratte S, Bertani A, Flecher X, Aubaniac JM. Long-term results with a lateral unicompartmental knee replacement. *Clin Orthop Relat Res* 2008;466:2686–93. <http://dx.doi.org/10.1007/s11999-008-0351-z>.
- [121] Berend KR, Kocczun MC, George JW, Lombardi AV. Lateral unicompartmental knee arthroplasty through a lateral parapatellar approach has high early survivorship. *Clin Orthop Relat Res* 2012;470:77–83. <http://dx.doi.org/10.1007/s11999-011-2005-9>.
- [122] Demange MK, Von Keudell A, Probst C, Yoshioka H, Gomoll AH. Patient-specific implants for lateral unicompartmental knee arthroplasty. *Int Orthop* 2015; 1519–26. <http://dx.doi.org/10.1007/s00264-015-2678-x>.
- [123] Lustig S, Lording T, Frank F, Debette C, Servien E, Neyret P. Progression of medial osteoarthritis and long term results of lateral unicompartmental arthroplasty: 10 to 18 year follow-up of 54 consecutive implants. *Knee* 2014;21:S26–32. [http://dx.doi.org/10.1016/S0968-0160\(14\)50006-3](http://dx.doi.org/10.1016/S0968-0160(14)50006-3).
- [124] Marson B, Prasad N, Jenkins R, Lewis M. Lateral unicompartmental knee replacements: early results from a District General Hospital. *Eur J Orthop Surg Traumatol* 2014;24:987–91. <http://dx.doi.org/10.1007/s00590-013-1277-z>.
- [125] Pandit H, Jenkins C, Beard DJ, Price AJ, Gill HS, Dodd CAF, et al. Mobile bearing dislocation in lateral unicompartmental knee replacement. *Knee* 2010;17:392–7. <http://dx.doi.org/10.1016/j.knee.2009.10.007>.
- [126] Pennington DW, Swienckowski JJ, Lutes VB, Drake GN. Lateral unicompartmental knee arthroplasty: survivorship and technical considerations at an average follow-up of 12.4 years. *J Arthroplasty* 2006;21:13–7. <http://dx.doi.org/10.1016/j.arth.2004.11.021>.
- [127] Sah AP, Scott RD. Lateral unicompartmental knee arthroplasty through a medial approach. Study with an average five-year follow-up. *J Bone Joint Surg Am* 2007;89:1948–54. <http://dx.doi.org/10.2106/JBJS.F.01457>.
- [128] Streit MR, Walker T, Bruckner T, Merle C, Kretzer JP, Clarius M, et al. Mobile-bearing lateral unicompartmental knee replacement with the Oxford domed tibial component: an independent series. *J Bone Joint Surg Br* 2012;94-B:1356–61. <http://dx.doi.org/10.1302/0301-620X.94B10.29119>.
- [129] Weston-Simons JS, Pandit H, Kendrick BJL, Jenkins C, Barker K, Dodd CAF, et al. The mid-term outcomes of the Oxford Domed Lateral unicompartmental knee replacement. *Bone Joint J* 2014;96(B):59–64. <http://dx.doi.org/10.1302/0301-620X.96B1.31630>.
- [130] Akhbari P, Malak T, Dawson-Bowling S, East D, Miles K, Butler-Manuel PA. The Avon patellofemoral joint replacement: mid-term prospective results from an independent centre. *Clin Orthop Surg* 2015;7:171. <http://dx.doi.org/10.4055/cios.2015.7.2.171>.
- [131] Al-Hadithy N, Patel R, Navadgi B, Deo S, Hollinghurst D, Satish V. Mid-term results of the FPV patellofemoral joint replacement. *Knee* 2014;21:138–41. <http://dx.doi.org/10.1016/j.knee.2013.08.010>.
- [132] Argenson JN, Flecher X, Parratte S, Aubaniac JM. Patellofemoral arthroplasty: an update. *Clin Orthop Relat Res* 2005;440:50–3. <http://dx.doi.org/10.1097/01.blo.0000187061.27573.70>.
- [133] Benazzo F, Rossi SMP, Ghiara M. Partial knee arthroplasty: patellofemoral arthroplasty and combined unicompartmental and patellofemoral arthroplasty implants – general considerations and indications, technique and clinical experience. *Knee* 2014;21:S43–6. [http://dx.doi.org/10.1016/S0968-0160\(14\)50009-9](http://dx.doi.org/10.1016/S0968-0160(14)50009-9).
- [134] Butler JE, Shannon R. Patellofemoral arthroplasty with a custom-fit femoral prosthesis. *Orthopedics* 2009;32:81.
- [135] Cartier P, Sanouiller J-L, Khefacha A. Long-term results with the first patellofemoral prosthesis. *Clin Orthop Relat Res* 2005;47:54. <http://dx.doi.org/10.1097/01.blo.0000171918.24998.d1>.
- [136] Charalambous CP, Abididin Z, Mills SP, Rogers S, Sutton P, Parkinson R. The low contact stress patellofemoral replacement: high early failure rate. *J Bone Joint Surg Br* 2011;93:484–9. <http://dx.doi.org/10.1302/0301-620X.93B4.25899>.
- [137] Dahm DL, Kalisvaart MM, Stuart MJ, Slettedahl SW. Patellofemoral arthroplasty: outcomes and factors associated with early progression of tibiofemoral arthritis. *Knee Surg Sports Traumatol Arthrosc* 2014;22:2554–9. <http://dx.doi.org/10.1007/s00167-014-3202-3>.
- [138] Feucht MJ, Cotic M, Beitzel K, Baldini JF, Meidinger G, Schöttle PB, et al. A matched-pair comparison of inlay and onlay trochlear designs for patellofemoral arthroplasty: no differences in clinical outcome but less progression of osteoarthritis with inlay designs. *Knee Surg Sports Traumatol Arthrosc* 2015. <http://dx.doi.org/10.1007/s00167-015-3733-2>.
- [139] Hernigou P, Caton J. Design, operative technique and ten-year results of the Hermes™ patellofemoral arthroplasty. *Int Orthop* 2014;38:437–42. <http://dx.doi.org/10.1007/s00264-013-2158-0>.
- [140] Hernigou P, Flouzat-Lachaniette CH, Delblond W, Duffiet P, Julian D. Computer-assisted navigation in patellofemoral arthroplasty: a new technique to improve rotational position of the trochlea. *HSS J* 2013;9:118–22. <http://dx.doi.org/10.1007/s11420-013-9328-x>.
- [141] Hofmann AA, McCandless JB, Shaeffer JF, Magee TH. Patellofemoral replacement: the third compartment. *Bone Joint J* 2013;95 B:124–8. <http://dx.doi.org/10.1053/j.sart.2008.11.014>.
- [142] Hoogervorst P, de Jong RJ, Hannink G, van Kampen A. A 21% conversion rate to total knee arthroplasty of a first-generation patellofemoral prosthesis at a mean follow-up of 9.7 years. *Int Orthop* 2015;39:1857–64. <http://dx.doi.org/10.1007/s00264-015-2941-1>.
- [143] Liow MHL, Goh GS-H, Tay DK-J, Chia S-L, Lo N-N, Yeo S-J. Obesity and the absence of trochlear dysplasia increase the risk of revision in patellofemoral arthroplasty. *Knee* 2015;9–15. <http://dx.doi.org/10.1016/j.knee.2015.05.009>.
- [144] Mohammed R, Jimulia T, Durve K, Bansal M, Green M, Learmonth D. Medium-term results of patellofemoral joint arthroplasty. *Acta Orthop Belg* 2008;74: 472–7.
- [145] Mont MA, Johnson AJ, Naziri Q, Kolisek FR, Leadbetter WB. Patellofemoral arthroplasty. 7-year mean follow-up. *J Arthroplasty* 2012;27:358–61. <http://dx.doi.org/10.1016/j.arth.2011.07.010>.
- [146] Morris MJ, Lombardi AV, Berend KR, Hurst JM, Adams JB. Clinical results of patellofemoral arthroplasty. *J Arthroplasty* 2013;28:199–201. <http://dx.doi.org/10.1016/j.arth.2013.05.012>.
- [147] Nicol SG, Loveridge JM, Weale AE, Ackroyd CE, Newman JH. Arthritis progression after patellofemoral joint replacement. *Knee* 2006;13:290–5. <http://dx.doi.org/10.1016/j.knee.2006.04.005>.
- [148] Odumenya M, Costa ML, Parsons N, Achten J, Dhillon M, Krikler SJ. The Avon patellofemoral joint replacement: five-year results from an independent centre. *J Bone Joint Surg Br* 2010;92:56–60. <http://dx.doi.org/10.1302/0301-620X.92B1.23135>.
- [149] Sarda P, Maheswaran S, Shetty A. Medium term results of Avon patellofemoral joint replacement. *Indian J Orthop* 2011;45:439. <http://dx.doi.org/10.4103/0019-5413.83761>.
- [150] Sisto DJ, Sarin VK. Custom patellofemoral arthroplasty of the knee. Surgical technique. *J Bone Joint Surg Am* 2007;89(Suppl. 2):214–25. <http://dx.doi.org/10.2106/JBJS.G.00186>.

- [151] Van Wagenberg JMF, Speigner B, Gosens T, De Waal Malefijt J. Midterm clinical results of the Autocentric II patellofemoral prosthesis. *Int Orthop* 2009;33:1603–8. <http://dx.doi.org/10.1007/s00264-009-0719-z>.
- [152] Williams DP, Pandit HG, Athanasou NA, Murray DW, Gibbons CLMH. Early revisions of the Femoro-Patella Vialla joint replacement. *Bone Joint J* 2013;95 B:793–7. <http://dx.doi.org/10.1302/0301-620X.95B6.31355>.
- [153] Yadav B, Shaw D, Radcliffe G, Dacheppalli S, Kluge W. Mobile-bearing, congruent patellofemoral prosthesis: short-term results. *J Orthop Surg (Hong Kong)* 2012;20:348–52.
- [154] van der List JP, Chawla H, Villa JC, Zuiderbaan HA, Pearle AD. Early functional outcome after lateral UKA is sensitive to postoperative lower limb alignment. *Knee Surg Sports Traumatol Arthrosc* 2015. <http://dx.doi.org/10.1007/s00167-015-3877-0>.
- [155] Zuiderbaan HA, van der List JP, Chawla H, Khamaisy S, Thein R, Pearle AD. Predictors of subjective outcome after medial unicompartmental knee arthroplasty. *J Arthroplasty* 2016. <http://dx.doi.org/10.1016/j.arth.2015.12.038>.
- [156] Hermigou P, Deschamps G. Alignment influences wear in the knee after medial unicompartmental arthroplasty. *Clin Orthop Relat Res* 2004;161–5.
- [157] van der List JP, Zuiderbaan HA, Pearle AD. Why do medial unicompartmental knee arthroplasties fail today? *J Arthroplasty* 2015;1–6. <http://dx.doi.org/10.1016/j.arth.2015.11.030>.
- [158] Pandit HG, Campi S, Hamilton TW, Dada OD, Pollalis S, Jenkins C, et al. Five-year experience of cementless Oxford unicompartmental knee replacement. *Knee Surg Sports Traumatol Arthrosc* 2015;1–9. <http://dx.doi.org/10.1007/s00167-015-3879-y>.
- [159] Liddle AD, Pandit H, Judge A, Murray DW. Effect of surgical caseload on revision rate following total and unicompartmental knee replacement. *J Bone Joint Surg Am* 2016;98:1–8. <http://dx.doi.org/10.2106/JBJS.N.00487>.
- [160] Goodfellow JW, O'Connor JJ, Murray DW. A critique of revision rate as an outcome measure: re-interpretation of knee joint registry data. *J Bone Joint Surg Br* 2010;92:1628–31. <http://dx.doi.org/10.1302/0301-620X.92B12.25193>.
- [161] Citak M, Suero EM, Citak M, Dunbar NJ, Branch SH, Condit MA, et al. Unicompartmental knee arthroplasty: is robotic technology more accurate than conventional technique? *Knee* 2013;20:268–71. <http://dx.doi.org/10.1016/j.knee.2012.11.001>.
- [162] Moon Y-W, Ha C, Do K-H, Kim C-Y, Han J-H, Na S-E, et al. Comparison of robot-assisted and conventional total knee arthroplasty: a controlled cadaver study using multiparameter quantitative three-dimensional CT assessment of alignment. *Comput Aided Surg* 2012;17:86–95. <http://dx.doi.org/10.3109/10929088.2012.654408>.
- [163] Chawla H, Ghomrawi HM, van der List JP, Eggman AA, Zuiderbaan HA, Pearle AD. Establishing age-specific cost-effective annual revision rates for unicompartmental knee arthroplasty: a meta-analysis. *J Arthroplasty* 2016.
- [164] Badawy M, Espehaug B, Indrekvam K, Havelin LI, Furnes O. Higher revision risk for unicompartmental knee arthroplasty in low-volume hospitals. *Acta Orthop* 2014;85:342–7. <http://dx.doi.org/10.3109/17453674.2014.920990>.
- [165] List JPVD, McDonald LS, Pearle AD. Systematic review of medial versus lateral survivorship in unicompartmental knee arthroplasty. *Knee* 2016;22:454–60. <http://dx.doi.org/10.1016/j.knee.2015.09.011>.
- [166] Heyse TJ, Khefacha A, Peersman G, Cartier P. Survivorship of UKA in the middle-aged. *Knee* 2012;19:585–91. <http://dx.doi.org/10.1016/j.knee.2011.09.002>.
- [167] Lonner JH. Patellofemoral arthroplasty: pros, cons, and design considerations. *Clin Orthop Relat Res* 2004;158–65.
- [168] Lustig S. Patellofemoral arthroplasty. *Orthop Traumatol Surg Res* 2014;100:S35–43. <http://dx.doi.org/10.1016/j.otsr.2013.06.013>.
- [169] Kozinn SC. Current Concept Review Unocpndylar Knee Arthroplasty 1989;71:145–50.
- [170] Argenson J-NA, Flecher X, Parratte S, Aubaniac J-M. Patellofemoral arthroplasty. *Clin Orthop Relat Res* 2005;440:50–3. <http://dx.doi.org/10.1097/01.blo.0000187061.27573.70>.
- [171] De Cloedt P, Legaye J, Lokietek W. Femoro-patellar prosthesis. A retrospective study of 45 consecutive cases with a follow-up of 3–12 years. *Acta rthop Belg* 1999;65:170–5.
- [172] Lonner JH. Patellofemoral arthroplasty: the impact of design on outcomes. *Orthop Clin North Am* 2008;39:347–54. <http://dx.doi.org/10.1016/j.jocl.2008.02.002>.
- [173] Cossey AJ, Spriggins AJ. Computer-assisted patellofemoral arthroplasty. A mechanism for optimizing rotation. *J Arthroplasty* 2006;21:420–7. <http://dx.doi.org/10.1016/j.arth.2005.08.010>.
- [174] Leadbetter WB. Patellofemoral arthroplasty in the treatment of patellofemoral arthritis: rationale and outcomes in younger patients. *Orthop Clin North Am* 2008;39:363–80. <http://dx.doi.org/10.1016/j.jocl.2008.04.001>.
- [175] Bull AMJ, Kessler O, Alam M, Amis AA. Changes in knee kinematics reflect the articular geometry after arthroplasty. *Clin Orthop Relat Res* 2008;466:2491–9. <http://dx.doi.org/10.1007/s11999-008-0440-z>.
- [176] Noble PC, Gordon MJ, Weiss JM, Reddix RN, Condit MA, Mathis KB. Does total knee replacement restore normal knee function? *Clin Orthop Relat Res* 2005;157–65. <http://dx.doi.org/10.1097/01.blo.0000150130.03519.fb>.
- [177] Ghomrawi HM, Eggman AA, Pearle AD. Effect of age on cost-effectiveness of unicompartmental knee arthroplasty compared with total knee arthroplasty in the U.S. *J Bone Joint Surg Am* 2015;97:396–402. <http://dx.doi.org/10.2106/JBJS.N.00169>.
- [178] Slover J, Espehaug B, Havelin LI, Engesaeter LB, Furnes O, Tomek I, et al. Cost-effectiveness of unicompartmental and total knee arthroplasty in elderly low-demand patients. A Markov decision analysis. *J Bone Joint Surg Am* 2006;88:2348–55. <http://dx.doi.org/10.2106/JBJS.E.01033>.
- [179] Nwachukwu BU, Mclawhorn AS, Simon MS, Hamid KS, Demetracopoulos CA, Deland JT, et al. Management of end-stage ankle arthritis: cost-utility analysis using direct and indirect costs; 2015 1159–72.
- [180] van Jonbergen H-PW, Poolman RW, van Kampen A. Isolated patellofemoral osteoarthritis. *Acta Orthop* 2010;81:199–205. <http://dx.doi.org/10.3109/17453671003628756>.
- [181] van der List J, Chawla H, Villa JC, Pearle AD. Why do patellofemoral arthroplasties fail today? *Knee* 2015. <http://dx.doi.org/10.1017/CBO9781107415324.004>.